PubMed	
--------	--

Format: Abstract

V

J Biol Regul Homeost Agents. 2018 Mar-Apr;32(2):341-344.

Correlation between childhood eczema and specific IgG antibody level.

Liu Y¹, Yan H², Shao F³, Li QH², Cui M².

Author information

Abstract

Eczema, a common pediatric dermatosis with unclear pathogenesis, can seriously affect the life quality of children due to its recurrence and long course. Recent study has found that food specific **IgG** (slgG) might be involved in the course of eczema. To analyze the correlation between childhood eczema and slgG and evaluate the role of avoiding taking intolerance food in the treatment of childhood eczema, this study enrolled 216 children with eczema who were admitted to the Taian Maternal and Child Health Care Hospital, Shandong, China, between August 2014 and October 2015. They were divided into an eczema group (N = 140) and an allergy group (N = 76). Eighty healthy children who were admitted to the Department of Children Healthcare in the same period were selected as a control group. Enzymelinked immuno sorbent assay (ELISA) was used to detect the serum slgG level. The result showed that the slgG positive rates of children in the eczema group and allergy group were 91.4% and 93.4%, respectively, and the difference had no statistical significance (P > 0.05). However, the slgG positive rates of children in the eczema group and allergy group were significantly higher than that in the control group (P less than 0.05). Milk and eggs were the major allergy-causing **food** for children with slgG positive rates higher than 70%. The slgG test results revealed that eggs had the highest allergenicity, followed by milk, tomatoes and soybeans, and pork was not highly sensitive. Therefore, it can be concluded that slgG positive rate of children with eczema is high, and examination of **food** slgG antibody in serum is valuable in the diagnosis and treatment of childhood eczema.

PMID: 29685016 [Indexed for MEDLINE]

Publication type, MeSH terms, Substance